

INVARIANTS OF COMPLEX STRUCTURES
ON NILMANIFOLDS

EDWIN ALEJANDRO RODRÍGUEZ VALENCIA

ABSTRACT. Let (N, J) be a simply connected $2n$ -dimensional nilpotent Lie group endowed with an invariant complex structure. We define a left invariant Riemannian metric on N compatible with J to be *minimal*, if it minimizes the norm of the invariant part of the Ricci tensor among all compatible metrics with the same scalar curvature. In [7], J. Lauret proved that minimal metrics (if any) are unique up to isometry and scaling. This uniqueness allows us to distinguish two complex structures with Riemannian data, giving rise to a great deal of invariants.

We show how to use a Riemannian invariant: the eigenvalues of the Ricci operator, polynomial invariants and discrete invariants to give an alternative proof of the pairwise non-isomorphism between the structures which have appeared in the classification of abelian complex structures on 6-dimensional nilpotent Lie algebras given in [1]. We also present some continuous families in dimension 8.

1. INTRODUCTION

Let N be a real $2n$ -dimensional nilpotent Lie group with Lie algebra \mathfrak{n} , whose Lie bracket will be denoted by $\mu: \mathfrak{n} \times \mathfrak{n} \rightarrow \mathfrak{n}$. An *invariant complex structure* on N is defined by a map $J: \mathfrak{n} \rightarrow \mathfrak{n}$ satisfying $J^2 = -I$ and the integrability condition

$$(1) \quad \mu(JX, JY) = \mu(X, Y) + J\mu(JX, Y) + J\mu(X, JY), \quad \forall X, Y \in \mathfrak{n}.$$

By left translating J , one obtains a complex manifold (N, J) , as well as compact complex manifolds $(N/\Gamma, J)$ if N admits cocompact discrete subgroups Γ , which are usually called *nilmanifolds* and play an important role in complex geometry.

The automorphism group $\text{Aut}(\mathfrak{n})$ acts by conjugation on the set of all invariant complex structures on \mathfrak{n} , and hence two such structures are considered to be *equivalent* if they belong to the same conjugation class. The lack of invariants makes the classification of invariant complex structures a difficult task. This has

2010 *Mathematics Subject Classification*: primary 53C15; secondary 32Q60, 53C30, 22E25, 37J15.

Key words and phrases: complex, nilmanifolds, nilpotent Lie groups, minimal metrics, Pfaffian forms.

Fully supported by a CONICET fellowship (Argentina).

Received September 14, 2014. Editor J. Slovák.

DOI: 10.5817/AM2015-1-27

only been achieved in dimension ≤ 6 in the nilpotent case in [2], and for any 6-dimensional Lie algebra in the abelian case in [1].

Our aim in this paper is to use two different invariants (namely, minimal metrics and Pfaffian forms, see below), to give an alternative proof of the non-equivalence between any two abelian complex structures on nilpotent Lie algebras of dimension 6 obtained in the classification list given in [1, Theorem 3.4.]. Along the way, we prove that any such structure, excepting only one, does admit a minimal metric. As another application of the invariants, we give in Section 5 many families depending on one, two and three parameters of abelian complex structures on 8-dimensional 2-step nilpotent Lie algebras, showing that a full classification could be really difficult in dimension 8.

1.1. Minimal metrics.

A left invariant metric which is *compatible* with (N, J) , also called a *Hermitian metric*, is determined by an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} such that

$$\langle JX, JY \rangle = \langle X, Y \rangle, \quad \forall X, Y \in \mathfrak{n}.$$

We consider

$$\text{Ric}^c_{\langle \cdot, \cdot \rangle} := \frac{1}{2} (\text{Ric}_{\langle \cdot, \cdot \rangle} - J \text{Ric}_{\langle \cdot, \cdot \rangle} J),$$

the complexified part of the Ricci operator $\text{Ric}_{\langle \cdot, \cdot \rangle}$ of the Hermitian manifold $(N, J, \langle \cdot, \cdot \rangle)$, and the corresponding $(1, 1)$ -component of the Ricci tensor $\text{ric}^c_{\langle \cdot, \cdot \rangle} := \langle \text{Ric}^c_{\langle \cdot, \cdot \rangle} \cdot, \cdot \rangle$.

A compatible metric $\langle \cdot, \cdot \rangle$ on (N, J) is called *minimal* if

$$\text{tr} (\text{Ric}^c_{\langle \cdot, \cdot \rangle})^2 = \min \{ \text{tr} (\text{Ric}^c_{\langle \cdot, \cdot \rangle'})^2 : \text{sc}(\langle \cdot, \cdot \rangle') = \text{sc}(\langle \cdot, \cdot \rangle) \},$$

where $\langle \cdot, \cdot \rangle'$ runs over all compatible metrics on (N, J) and $\text{sc}(\langle \cdot, \cdot \rangle) = \text{tr} \text{Ric}^c_{\langle \cdot, \cdot \rangle}$ is the scalar curvature. In [7], the following conditions on $\langle \cdot, \cdot \rangle$ are proved to be equivalent to minimality, showing that such metrics are special from many other points of view:

- (i) The solution $\langle \cdot, \cdot \rangle_t$ with initial value $\langle \cdot, \cdot \rangle_0 = \langle \cdot, \cdot \rangle$ to the evolution equation

$$\frac{d}{dt} \langle \cdot, \cdot \rangle_t = -2 \text{ric}^c_{\langle \cdot, \cdot \rangle_t},$$

is self-similar, in the sense that $\langle \cdot, \cdot \rangle_t = c_t \varphi_t^* \langle \cdot, \cdot \rangle$ for some $c_t > 0$ and one-parameter group of automorphisms φ_t of N .

- (ii) There exist a vector field X on N and $c \in \mathbb{R}$ such that

$$\text{ric}^c_{\langle \cdot, \cdot \rangle} = c \langle \cdot, \cdot \rangle + L_X \langle \cdot, \cdot \rangle,$$

where $L_X \langle \cdot, \cdot \rangle$ denotes the usual Lie derivative. In analogy with the well-known concept in Ricci flow theory, one may call $\langle \cdot, \cdot \rangle$ a $(1, 1)$ -Ricci soliton.

- (iii) $\text{Ric}^c_{\langle \cdot, \cdot \rangle} = cI + D$ for some $c \in \mathbb{R}$ and $D \in \text{Der}(\mathfrak{n})$.

The uniqueness up to isometric isomorphism and scaling of a minimal metric on a given (N, J) was also proved in [7], and can be used to obtain invariants in the following way. If $(N, J_1, \langle \cdot, \cdot \rangle_1)$ and $(N, J_2, \langle \cdot, \cdot \rangle_2)$ are minimal and J_1 is equivalent to J_2 , then they must be conjugate via an automorphism which is an isometry

between $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$. This provides us with a lot of invariants, namely the Riemannian geometry invariants including all different kind of curvatures.

1.2. Pfaffian forms.

Consider a real vector space \mathfrak{n} and fix a direct sum decomposition

$$\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2, \quad \dim \mathfrak{n}_1 = m, \quad \dim \mathfrak{n}_2 = n.$$

Every 2-step nilpotent Lie algebra of dimension $m + n$ with derived algebra of dimension $\leq n$ can be represented by a bilinear skew-symmetric map

$$\mu: \mathfrak{n}_1 \times \mathfrak{n}_1 \longrightarrow \mathfrak{n}_2.$$

For a given inner product $\langle \cdot, \cdot \rangle$ on $\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2$ (with $\mathfrak{n}_1 \perp \mathfrak{n}_2$), one can encode the structural constants of μ in a map $J_\mu: \mathfrak{n}_2 \longrightarrow \mathfrak{so}(\mathfrak{n}_1)$ defined by

$$\langle J_\mu(Z)X, Y \rangle = \langle \mu(X, Y), Z \rangle, \quad \forall X, Y \in \mathfrak{n}_1, Z \in \mathfrak{n}_2.$$

There is a nice and useful isomorphism invariant for 2-step algebras (with m even) called the *Pfaffian form*, which is the projective equivalence class of the homogeneous polynomial f_μ of degree $m/2$ in n variables defined by

$$f_\mu(Z)^2 = \det J_\mu(Z), \quad \forall Z \in \mathfrak{n}_2,$$

for each μ of type (n, m) (see Section 3.2).

For each $\mu \in V_{n,m} := \Lambda^2 \mathfrak{n}_1^* \otimes \mathfrak{n}_2$, let N_μ denote the simply connected nilpotent Lie group with Lie algebra (\mathfrak{n}, μ) . We prove that if two complex nilmanifolds (N_μ, J) and (N_λ, J) are *holomorphically isomorphic*, then $f_\lambda \in \mathbb{R}_{>0} GL_q(\mathbb{C}) \cdot f_\mu$, with $n = 2q$ (see Proposition 3.12). This will allow us to use the existence of minimal metrics to distinguish complex nilmanifolds by means of invariants of forms.

Acknowledgement. This research is part of the Ph.D. thesis (Universidad Nacional de Córdoba) by the author. I am grateful to my advisor Dr. Jorge Lauret for his invaluable help during the preparation of the paper.

2. PRELIMINARIES

In this section, we recall basic notions on complex structures on nilmanifolds and their Hermitian metrics.

Let N be a simply connected $2n$ -dimensional nilpotent Lie group with Lie algebra \mathfrak{n} , whose Lie bracket will be denoted by $\mu: \mathfrak{n} \times \mathfrak{n} \rightarrow \mathfrak{n}$. An invariant *complex structure* on N is defined by a map $J: \mathfrak{n} \rightarrow \mathfrak{n}$ satisfying $J^2 = -I$ and such that

$$\mu(JX, JY) = \mu(X, Y) + J\mu(JX, Y) + J\mu(X, JY), \quad \forall X, Y \in \mathfrak{n}.$$

We say that J is *abelian* if the following condition holds:

$$\mu(JX, JY) = \mu(X, Y), \quad \forall X, Y \in \mathfrak{n}.$$

Definition 2.1. Two complex structures J_1 and J_2 on N are said to be *equivalent* if there exists an automorphism α of \mathfrak{n} satisfying $J_2 = \alpha J_1 \alpha^{-1}$. Two pairs (N_1, J_1) and (N_2, J_2) are *holomorphically isomorphic* if there exists a Lie algebra isomorphism $\alpha: \mathfrak{n}_1 \rightarrow \mathfrak{n}_2$ such that $J_2 = \alpha J_1 \alpha^{-1}$.

We fix a $2n$ -dimensional real vector space \mathfrak{n} , and consider as a parameter space for the set of all real nilpotent Lie algebras of a given dimension $2n$, the algebraic subset

$$\mathcal{N} := \{\mu \in V : \mu \text{ satisfies Jacobi and is nilpotent}\},$$

where $V := \Lambda^2 \mathfrak{n}^* \otimes \mathfrak{n}$ is the vector space of all skew-symmetric bilinear maps from $\mathfrak{n} \times \mathfrak{n}$ to \mathfrak{n} . Recall that any inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} determines an inner product on V , also denoted by $\langle \cdot, \cdot \rangle$, as follows: if $\{e_i\}$ is an orthonormal basis of \mathfrak{n} ,

$$\begin{aligned} \langle \mu, \lambda \rangle &:= \sum_{i,j} \langle \mu(e_i, e_j), \lambda(e_i, e_j) \rangle \\ (2) \quad &= \sum_{i,j,k} \langle \mu(e_i, e_j), e_k \rangle \langle \lambda(e_i, e_j), e_k \rangle. \end{aligned}$$

For each $\mu \in \mathcal{N}$, let N_μ denote the simply connected nilpotent Lie group with Lie algebra (\mathfrak{n}, μ) . We now fix a map $J: \mathfrak{n} \rightarrow \mathfrak{n}$ such that $J^2 = -I$. The corresponding Lie group

$$GL_n(\mathbb{C}) = \{g \in GL_{2n}(\mathbb{R}) : gJ = Jg\}$$

acts naturally on V by $g \cdot \mu(\cdot, \cdot) = g\mu(g^{-1}\cdot, g^{-1}\cdot)$, leaving \mathcal{N} invariant, as well as the algebraic subset $\mathcal{N}_J \subset \mathcal{N}$ given by

$$\mathcal{N}_J := \{\mu \in \mathcal{N} : \mu \text{ satisfies (1)}\}.$$

We can identify each $\mu \in \mathcal{N}_J$ with a *complex nilmanifold* as follows:

$$(3) \quad \mu \leftrightarrow (N_\mu, J).$$

Proposition 2.2. *Two complex nilmanifolds μ and λ are holomorphically isomorphic if and only if $\lambda \in GL_n(\mathbb{C}) \cdot \mu$.*

Proof. If we suppose that (N_μ, J) and (N_λ, J) are holomorphically isomorphic, then there exists a Lie algebra isomorphism $g^{-1}: (\mathfrak{n}, \lambda) \mapsto (\mathfrak{n}, \mu)$ such that $J = gJg^{-1}$. Hence, $\lambda = g \cdot \mu$ and $g \in GL_n(\mathbb{C})$ (taking their matrix representation). \square

A left invariant Riemannian metric on N is said to be *compatible* with a complex structure J on N if it is defined by an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} such that

$$\langle JX, JY \rangle = \langle X, Y \rangle, \quad \forall X, Y \in \mathfrak{n},$$

that is, J is orthogonal with respect to $\langle \cdot, \cdot \rangle$. We denote by $\mathcal{C} = \mathcal{C}(N, J)$ the set of all left invariant metrics on N compatible with J .

Definition 2.3. Two triples $(N_1, J_1, \langle \cdot, \cdot \rangle)$ and $(N_2, J_2, \langle \cdot, \cdot \rangle')$, with $\langle \cdot, \cdot \rangle \in \mathcal{C}(N_1, J_1)$ and $\langle \cdot, \cdot \rangle' \in \mathcal{C}(N_2, J_2)$, are said to be *isometric isomorphic* if there exists a Lie algebra isomorphism $\varphi: \mathfrak{n}_1 \rightarrow \mathfrak{n}_2$ such that $J_2 = \varphi J_1 \varphi^{-1}$ and $\langle \cdot, \cdot \rangle' = \langle \varphi^{-1}\cdot, \varphi^{-1}\cdot \rangle$.

We now identify each $\mu \in \mathcal{N}_J$ with a *Hermitian nilmanifold* in the following way:

$$(4) \quad \mu \leftrightarrow (N_\mu, J, \langle \cdot, \cdot \rangle),$$

where $\langle \cdot, \cdot \rangle$ is a fixed inner product on \mathfrak{n} compatible with J . Therefore, each $\mu \in \mathcal{N}_J$ can be viewed in this way as a *Hermitian metric* compatible with (N_μ, J) , and two metrics μ, λ are compatible with the same complex structure if and only if they

live in the same $GL_n(\mathbb{C})$ -orbit. Indeed, each $g \in GL_n(\mathbb{C})$ determines a Riemannian isometry preserving the complex structure

$$(5) \quad (N_{g \cdot \mu}, J, \langle \cdot, \cdot \rangle) \rightarrow (N_\mu, J, \langle g \cdot, g \cdot \rangle)$$

by exponentiating the Lie algebra isomorphism $g^{-1}: (\mathfrak{n}, g \cdot \mu) \mapsto (\mathfrak{n}, \mu)$. We then have the identification $GL_n(\mathbb{C}) \cdot \mu = \mathcal{C}(N_\mu, J)$, for any $\mu \in \mathcal{N}_J$.

3. INVARIANTS

We now discuss the problem of distinguishing two complex nilmanifolds up to holomorphic isomorphism, by considering different types of invariants.

3.1. Minimal metrics.

In [7], J. Lauret showed how to use the complexified part of the Ricci operator of a nilpotent Lie group given, to determinate the existence of compatible *minimal* metrics with an invariant geometric structure on the Lie group. Furthermore, he proved that these metrics (if any) are unique up to isometry and scaling. This property allows us to distinguish two geometric structure with invariants coming from Riemannian geometric. In this section, we will be apply these results to the complex case and use the identifications (3) and (4) to rewrite them in terms of data arising from the Lie algebra; this will be the basis of our method: fix a complex structure and move the bracket. This method is explained in a more detailed way in Section 4 in the 6-dimensional case.

The following theorem was obtained by using strong results from geometric invariant theory, mainly related to the moment map of a real representation of a real reductive Lie group.

Theorem 3.1 ([7]). *Let $F: \mathcal{N}_J \rightarrow \mathbb{R}$ be defined by $F(\mu) := \text{tr}(\text{Ric}_\mu^c)^2 / \|\mu\|^4$, where Ric_μ^c is the orthogonal projection of the Ricci operator Ric_μ of the Riemannian manifold $(N_\mu, \langle \cdot, \cdot \rangle)$ onto the space of symmetric maps of $(\mathfrak{n}, \langle \cdot, \cdot \rangle)$ which commute with J . Then for $\mu \in \mathcal{N}_J$, the following conditions are equivalent:*

- (i) μ is a critical point of F .
- (ii) $F|_{GL_n(\mathbb{C}) \cdot \mu}$ attains its minimum value at μ .
- (iii) $\text{Ric}_\mu^c = cI + D$ for some $c \in \mathbb{R}$, $D \in \text{Der}(\mathfrak{n})$.

Moreover, all the other critical points of F in the orbit $GL_n(\mathbb{C}) \cdot \mu$ lie in $\mathbb{R}^*U(n) \cdot \mu$.

A complex nilmanifold μ is said to be *minimal* if it satisfies any of the conditions in the previous theorem.

Corollary 3.2. *Two minimal complex nilmanifolds μ and λ are holomorphically isomorphic if and only if $\lambda \in \mathbb{R}^*U(n) \cdot \mu$.*

Let $(N, J, \langle \cdot, \cdot \rangle)$ be a Hermitian nilmanifold, i.e. J is an invariant complex structure on N and $\langle \cdot, \cdot \rangle \in \mathcal{C}(N, J)$.

Definition 3.3. Let $\text{Ric}_{\langle \cdot, \cdot \rangle}$ be the Ricci operator of $(N, \langle \cdot, \cdot \rangle)$. The *Hermitian Ricci operator* is given by

$$\text{Ric}_{\langle \cdot, \cdot \rangle}^c := \frac{1}{2} (\text{Ric}_{\langle \cdot, \cdot \rangle} - J \text{Ric}_{\langle \cdot, \cdot \rangle} J) .$$

A metric $\langle \cdot, \cdot \rangle \in \mathcal{C}$ is called *minimal* if it minimizes the functional $\text{tr}(\text{Ric}^c_{\langle \cdot, \cdot \rangle})^2$ on the set of all compatible metrics with the same scalar curvature. We now rewrite Theorem 3.1 in geometric terms, by using the identification (4).

Theorem 3.4 ([7]). *For $\langle \cdot, \cdot \rangle \in \mathcal{C}$, the following conditions are equivalent:*

- (i) $\langle \cdot, \cdot \rangle$ is minimal.
- (ii) $\text{Ric}^c_{\langle \cdot, \cdot \rangle} = cI + D$ for some $c \in \mathbb{R}$, $D \in \text{Der}(\mathfrak{n})$.

Moreover, there is at most one compatible left invariant metric on (N, J) up to isometry (and scaling) satisfying any of the above conditions.

Let $\langle \cdot, \cdot \rangle \in \mathcal{C}$ be a minimal metric with $\text{Ric}^c_{\langle \cdot, \cdot \rangle} = cI + D$ for some $c \in \mathbb{R}$, $D \in \text{Der}(\mathfrak{n})$. We say that μ is of *type* $(k_1 < \dots < k_r; d_1, \dots, d_r)$ if $\{k_i\} \subset \mathbb{Z}_{\geq 0}$ are the eigenvalues of D with multiplicities $\{d_i\}$ respectively and $\text{gcd}(k_1, \dots, k_r) = 1$.

Corollary 3.5 ([7]). *Let J_1, J_2 be two complex structures on N , and assume that they admit minimal compatible metrics $\langle \cdot, \cdot \rangle$ and $\langle \cdot, \cdot \rangle'$, respectively. Then J_1 is equivalent to J_2 if and only if there exists $\varphi \in \text{Aut}(\mathfrak{n})$ and $c > 0$ such that $J_2 = \varphi J_1 \varphi^{-1}$ and*

$$\langle \varphi X, \varphi Y \rangle' = c \langle X, Y \rangle, \quad \forall X, Y \in \mathfrak{n}.$$

In particular, if J_1 and J_2 are equivalent, then their respective minimal compatible metrics are necessarily isometric up to scaling.

By (4) and (5), it is easy to see that two Hermitian nilmanifolds μ and λ are isometric (i.e. if $(N_\mu, J, \langle \cdot, \cdot \rangle)$ and $(N_\lambda, J, \langle \cdot, \cdot \rangle)$ are isometric isomorphic) if and only if they live in the same $U(n)$ -orbit. Corollary 3.5 and (4) imply the following result.

Corollary 3.6. *If μ is a minimal Hermitian metric, then $\mathbb{R}^*U(n) \cdot \mu$ parameterizes all minimal Hermitian metrics on (N_μ, J) .*

Example 3.7. For $t \in (0, 1]$, consider the 2-step nilpotent Lie algebra whose bracket is given by

$$\begin{aligned} \mu_t(e_1, e_2) &= \sqrt{t}e_5, & \mu_t(e_1, e_4) &= \frac{1}{\sqrt{t}}e_6, \\ \mu_t(e_2, e_3) &= -\frac{1}{\sqrt{t}}e_6, & \mu_t(e_3, e_4) &= -\sqrt{t}e_5. \end{aligned}$$

Let

$$J = \begin{bmatrix} 0 & -1 & & & & \\ 1 & 0 & & & & \\ & & 0 & -1 & & \\ & & 1 & 0 & & \\ & & & & 0 & -1 \\ & & & & 1 & 0 \end{bmatrix}, \quad \langle e_i, e_j \rangle = \delta_{ij}.$$

A straightforward verification shows that J is an abelian complex structure on N_{μ_t} for all t , $\langle \cdot, \cdot \rangle$ is compatible with (N_{μ_t}, J) , and the Ricci operator is given by

$$\text{Ric}_{\mu_t} = \begin{bmatrix} -\frac{1}{2} \left(\frac{t^2+1}{t} \right) I_4 & & & \\ & t & 0 & \\ & 0 & 1/t & \end{bmatrix}.$$

By definition, we have

$$\text{Ric}^c_{\mu_t} = \begin{bmatrix} -\left(\frac{t^2+1}{2t} \right) I_4 & & & \\ & \left(\frac{t^2+1}{2t} \right) I_2 & & \end{bmatrix} = \frac{t^2+1}{2t} \left(-3I + 2 \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix} \right),$$

and thus μ_t is minimal of type $(1 < 2; 4, 2)$ by Theorem 3.4. It follows from

$$\text{Ric}_{\mu_t}|_{\mathfrak{n}_2} = \begin{bmatrix} t & 0 \\ 0 & 1/t \end{bmatrix},$$

that the Hermitian nilmanifolds $\{(N_{\mu_t}, J, \langle \cdot, \cdot \rangle) : 0 < t \leq 1\}$ are pairwise non-isometric. Indeed, if there exists $c \in \mathbb{R}^*$ and $\varphi \in \text{U}(3) \subset \text{O}(6)$ such that $c\mu_s = \varphi \cdot \mu_t$ (see Corollary 3.6), then $\varphi = [\varphi^1 \ \varphi_2] \in \text{U}(2) \times \text{U}(1)$ (recall that it is of type (4,2)) and $c^2 \text{Ric}_{\mu_s}|_{\mathfrak{n}_2} = \varphi_2 \text{Ric}_{\mu_t}|_{\mathfrak{n}_2} \varphi_2^{-1}$, hence $c^2 \begin{bmatrix} s & \\ & 1/s \end{bmatrix} = \begin{bmatrix} t & \\ & 1/t \end{bmatrix}$. By taking quotients of their eigenvalues we deduce that $s^2 = t^2$ or $s^2 = 1/t^2$, which gives $s = t$ if $s, t \in (0, 1]$. We therefore obtain a curve $\{(N_{\mu_t}, J) : 0 < t \leq 1\}$ of pairwise non-isomorphic abelian complex nilpotent Lie groups, by the uniqueness in result Theorem 3.4 (see [6] for more examples).

From the above results, the problem of distinguishing two complex structures can be stated as follows: if we fix the nilpotent Lie group N then the $GL_{2n}(\mathbb{R})$ -invariants give us all possible complex structures on N (Definition 2.1), and the $\text{O}(2n)$ -invariants distinguish their respective minimal metrics (if any), up to scaling (Corollary 3.5). If we now fix a $2n$ -dimensional vector space and vary the brackets, the $GL_n(\mathbb{C})$ -invariants provide the possible compatible metrics with a given complex structure (see identification (4)), and the $\text{U}(n)$ -invariants their respective minimal metrics (if any), up to scaling (see Corollary 3.6). In the latter case, the above example shows how to use one of the Riemannian invariants: the eigenvalues of the Ricci operator. Since this is not always possible, in the next section we will introduce a new invariant applicable to 2-step nilpotent Lie algebras.

3.2. Pfaffian form.

With the purpose to differentiate Lie algebras, up to isomorphism, we assign to each one a unique homogeneous polynomial called the *Pfaffian form*, and by Proposition 3.10 we will use the known polynomial invariants to obtain curves or families of brackets in a vector space given. We follow the notation used in [8].

Let \mathfrak{n} be a real Lie algebra, with Lie bracket μ , and fix an inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{n} . For each $Z \in \mathfrak{n}$ consider the skew-symmetric \mathbb{R} -linear transformation $J_Z : \mathfrak{n} \rightarrow \mathfrak{n}$ defined by

$$(6) \quad \langle J_Z X, Y \rangle = \langle \mu(X, Y), Z \rangle, \quad \forall X, Y \in \mathfrak{n}.$$

If \mathfrak{n} and \mathfrak{n}' are two real Lie algebras and J, J' are the corresponding maps, relative to the inner products $\langle \cdot, \cdot \rangle$ and $\langle \cdot, \cdot \rangle'$ respectively, then it is easy to see that a linear map $B : \mathfrak{n} \rightarrow \mathfrak{n}'$ is a Lie algebra isomorphism if and only if

$$(7) \quad B^t J'_Z B = J_{B^t Z}, \quad \forall Z \in \mathfrak{n}',$$

where $B^t : \mathfrak{n}' \rightarrow \mathfrak{n}$ is given by $\langle B^t X, Y \rangle = \langle X, BY \rangle'$ for all $X \in \mathfrak{n}'$, $Y \in \mathfrak{n}$.

Assume now that \mathfrak{n} is 2-step nilpotent and the decomposition $\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2$ satisfies $\mathfrak{n}_2 = [\mathfrak{n}, \mathfrak{n}]$. If $\langle \mathfrak{n}_1, \mathfrak{n}_2 \rangle = 0$, then \mathfrak{n}_1 is J_Z -invariant for any Z and $J_Z = 0$ if and only if $Z \in \mathfrak{n}_1$. Under these conditions, the *Pfaffian form* $f : \mathfrak{n}_2 \rightarrow \mathbb{R}$ of \mathfrak{n} is defined by

$$f(Z) = \text{Pf}(J_Z|_{\mathfrak{n}_1}), \quad Z \in \mathfrak{n}_2,$$

Proposition 3.12. *Suppose that $\mathfrak{n} = \mathfrak{n}_1 \oplus \mathfrak{n}_2$, with $\dim \mathfrak{n}_1 = 2p$ and $\dim \mathfrak{n}_2 = 2q$, and $J\mathfrak{n}_i = \mathfrak{n}_i$. Assume $\mu, \lambda \in \Lambda^2 \mathfrak{n}_1^* \otimes \mathfrak{n}_2$ satisfy $\mu(\mathfrak{n}_1, \mathfrak{n}_1) = \lambda(\mathfrak{n}_1, \mathfrak{n}_1) = \mathfrak{n}_2$. If $\lambda \in GL_n(\mathbb{C}) \cdot \mu$ ($n=p+q$), then*

$$f(\lambda) \in \mathbb{R}_{>0}GL_q(\mathbb{C}) \cdot f(\mu),$$

where $f(\mu), f(\lambda)$ are the Pfaffian forms of (\mathfrak{n}, μ) and (\mathfrak{n}, λ) , respectively.

Proof. Let $\mathfrak{h} := (\mathfrak{n}, \mu)$, $\mathfrak{h}' := (\mathfrak{n}, \lambda)$ and J_μ, J_λ the corresponding maps, relative to the inner products on \mathfrak{n} (see (6)). Suppose that $g \cdot \mu = \lambda$ with $g \in GL_n(\mathbb{C})$ (i.e. $g \in GL_{2n}(\mathbb{R})$, $gJ = Jg$). By assumption, $g = \begin{bmatrix} g_1 & \\ & g_2 \end{bmatrix} \in GL_p(\mathbb{C}) \times GL_q(\mathbb{C})$ and $g: \mathfrak{h} \rightarrow \mathfrak{h}'$ is a Lie algebra isomorphism satisfying $g\mathfrak{n}_1 = \mathfrak{n}_1$ and $g\mathfrak{n}_2 = \mathfrak{n}_2$. It follows from (7) that

$$g^t J_\lambda(Z)g = J_\mu(g^t Z), \quad \forall Z \in \mathfrak{n}_1,$$

and since the subspaces \mathfrak{n}_1 and \mathfrak{n}_2 are preserved by g y g^t we have that

$$f'(Z) = cf(g_2^t Z),$$

where $c^{-1} = \det g_1 > 0$ ($GL_p(\mathbb{C})$ is connected) and $g_2^t: \lambda(\mathfrak{n}_1, \mathfrak{n}_1) \rightarrow \mu(\mathfrak{n}_1, \mathfrak{n}_1)$. It is clear that $g_2^t \in GL_{2q}(\mathbb{R})$ and satisfies

$$\langle Jg_2^t Z, Y \rangle = \langle g_2^t Z, -JY \rangle = \langle Z, g_2(-JY) \rangle = \langle Z, -Jg_2 Y \rangle = \langle JZ, g_2 Y \rangle = \langle g_2^t JZ, Y \rangle.$$

Thus $g_2^t \in GL_q(\mathbb{C})$ and we conclude that $f(\lambda) \in \mathbb{R}_{>0}GL_q(\mathbb{C}) \cdot f(\mu)$. \square

We end this section with an example of two homogeneous polynomials that are projectively equivalent over \mathbb{R} but not over \mathbb{C} (in the sense of Proposition 3.12).

Example 3.13. In $\mathfrak{h}_5 \times \mathbb{R}$, define the Lie brackets μ^+ and μ^- by

$$\mu^\pm(e_1, e_2) = e_6, \quad \mu^\pm(e_3, e_4) = \pm e_6.$$

Consider the inner product $\langle e_i, e_j \rangle = \delta_{ij}$. If $Z = xe_6$, with $x \in \mathbb{R}$, then

$$J_Z^+|_{\mathfrak{n}_1} = \begin{bmatrix} 0 & -x \\ x & 0 \\ & 0 & -x \\ & x & 0 \end{bmatrix}, \quad J_Z^-|_{\mathfrak{n}_1} = \begin{bmatrix} 0 & -x \\ x & 0 \\ & 0 & x \\ & -x & 0 \end{bmatrix}.$$

Hence $f(\mu^+) = x^2$ and $f(\mu^-) = -x^2$. It follows that $f(\mu^-) \simeq_{\mathbb{R}} f(\mu^+)$ but

$$f(\mu^-) \notin \mathbb{R}_{>0}U(1) \cdot f(\mu^+).$$

Recall that $GL_1(\mathbb{C}) = \mathbb{R}_{>0}U(1)$.

4. MINIMAL METRICS ON 6-DIMENSIONAL ABELIAN COMPLEX NILMANIFOLDS

The classification of 6-dimensional nilpotent real Lie algebras admitting a complex structure was given in [11], and the abelian case in [3]. Lately, A. Andrada, M.L. Barberis and I.G. Dotti in [1] gave a classification of all 6-dimensional Lie algebras admitting an abelian complex structure; furthermore, they give a parametrization, on each Lie algebra, of the space of abelian structures up to holomorphic isomorphism. In particular, there are three nilpotent Lie algebras carrying curves of non-equivalent structures. Based on this parametrization, we study the existence of minimal metrics on each of these complex nilmanifolds (see Theorem 4.4), and provide an alternative proof of the pairwise non-isomorphism between the structures.

The classification in [1] fix the Lie algebra and varies the complex structure. For example, on the Lie algebra $\mathfrak{h}_3 \times \mathfrak{h}_3$ they found the curve J_s of abelian complex structures defined by $J_s e_1 = e_2$, $J_s e_3 = e_4$, $J_s e_5 = s e_5 + e_6$, $s \in \mathbb{R}$, and fix the bracket $[e_1, e_2] = e_5$, $[e_3, e_4] = e_6$. We now fix the complex structure and varies the bracket as follows.

For $\mathfrak{n} = \mathfrak{v}_1 \oplus \mathfrak{v}_2$, with $\mathfrak{v}_1 = \mathbb{R}^4$ and $\mathfrak{v}_2 = \mathbb{R}^2$, consider the vector space $\Lambda^2 \mathfrak{v}_1^* \otimes \mathfrak{v}_2$ of all skew symmetric bilinear maps $\mu: \mathfrak{v}_1 \times \mathfrak{v}_1 \rightarrow \mathfrak{v}_2$. Any 6-dimensional 2-step nilpotent Lie algebra with $\dim \mu(\mathfrak{n}, \mathfrak{n}) \leq 2$ can be modelled in this way. Fix a basis of \mathfrak{n} , say $\{e_1, \dots, e_6\}$, such that $\mathfrak{v}_1 = \langle e_1, \dots, e_4 \rangle_{\mathbb{R}}$, $\mathfrak{v}_2 = \langle e_5, e_6 \rangle_{\mathbb{R}}$. The complex structure and the compatible metric will be always defined by

$$(9) \quad J := \begin{bmatrix} 0 & -1 & & & & \\ 1 & 0 & & & & \\ & & 0 & -1 & & \\ & & 1 & 0 & & \\ & & & & 0 & -1 \\ & & & & 1 & 0 \end{bmatrix}, \quad \langle e_i, e_j \rangle := \delta_{ij}.$$

Proposition 4.1. *Let $(N_{\tilde{\mu}}, \tilde{J})$ be a complex nilmanifold, with $\tilde{\mu} \in \Lambda^2 \mathfrak{v}_1^* \otimes \mathfrak{v}_2$. If there exists $g \in GL_6(\mathbb{R})$ such that $g\tilde{J}g^{-1} = J$, then $(N_{\tilde{\mu}}, \tilde{J})$ and $(N_{g\tilde{\mu}}, J)$ are holomorphically isomorphic.*

Returning to the above example, by choosing

$$g = \begin{bmatrix} 1 & 0 & & & & \\ 0 & 1 & & & & \\ & & 1 & 0 & & \\ & & 0 & 1 & & \\ & & & & 1 & -s \\ & & & & 0 & 1 \end{bmatrix},$$

we have $gJ_s g^{-1} = J$, and therefore $(N_{[\cdot, \cdot]}, J_s)$ and (N_{μ_3}, J) are holomorphically isomorphic by Proposition 4.1, where now the bracket is given by $\mu_3(e_1, e_2) = e_5$ and $\mu_3(e_3, e_4) = -s e_5 + e_6$ with $s \in \mathbb{R}$. By arguing as above for each item in [1, Theorem 3.4.], we have obtained Table 1.

Remark 4.2. In the classification given in [1], they incorrectly claim that the curves of structures J_t^1 and J_t^2 on \mathfrak{n}_4 are non-equivalent (see a corrected version at arXiv:0908.3213). Indeed, the matrix g defined in (10) is an automorphism of \mathfrak{n}_4 and $gJ_t^1 g^{-1} = J_t^2$, hence J_t^1 and J_t^2 are equivalent. Note that in Table 1 only appears a ‘curve’ (it is proved below) of brackets on \mathfrak{n}_4 , which is due to the following proposition and Theorem 4.4. The brackets $\mu_4^{1,t}$ and $\mu_4^{2,t}$ are obtained from the curves of structures J_t^1 and J_t^2 , respectively.

Proposition 4.3. $\mu_4^{2,t} \in U(2) \times U(1) \cdot \mu_4^{1,t}$ for all $t \in (0, 1]$, where the brackets $\mu_4^{1,t}, \mu_4^{2,t}$ on \mathfrak{n}_4 are given by

$$\begin{aligned} \mu_4^{1,t}(e_1, e_2) &= \sqrt{t} e_5, & \mu_4^{1,t}(e_1, e_4) &= \frac{1}{\sqrt{t}} e_6, & \mu_4^{2,t}(e_1, e_3) &= \sqrt{t} e_5, & \mu_4^{2,t}(e_2, e_4) &= \sqrt{t} e_5, \\ \mu_4^{1,t}(e_2, e_3) &= -\frac{1}{\sqrt{t}} e_6, & \mu_4^{1,t}(e_3, e_4) &= -\sqrt{t} e_5, & \mu_4^{2,t}(e_1, e_4) &= -\frac{1}{\sqrt{t}} e_6, & \mu_4^{2,t}(e_2, e_3) &= \frac{1}{\sqrt{t}} e_6. \end{aligned}$$

n	Bracket
$\mathfrak{n}_1 := \mathfrak{h}_3 \times \mathbb{R}^3$	$\mu_1(e_1, e_2) = e_6$
$\mathfrak{n}_2 := \mathfrak{h}_5 \times \mathbb{R}$	$\mu_2^\pm(e_1, e_2) = e_6, \mu_2^\pm(e_3, e_4) = \pm e_6$
$\mathfrak{n}_3 := \mathfrak{h}_3 \times \mathfrak{h}_3$	$\mu_3^s(e_1, e_2) = e_5, \mu_3^s(e_3, e_4) = -se_5 + e_6$ $s \in \mathbb{R}$
$\mathfrak{n}_4 := \mathfrak{h}_3(\mathbb{C})$	$\mu_4^t(e_1, e_2) = \sqrt{t}e_5, \mu_4^t(e_1, e_4) = \frac{1}{\sqrt{t}}e_6$ $\mu_4^t(e_2, e_3) = -\frac{1}{\sqrt{t}}e_6, \mu_4^t(e_3, e_4) = -\sqrt{t}e_5$ $t \in (0, 1]$
\mathfrak{n}_5	$\mu_5(e_1, e_2) = e_5, \mu_5(e_1, e_4) = -e_6$ $\mu_5(e_2, e_3) = e_6$
\mathfrak{n}_6	$\mu_6(e_1, e_2) = -e_3, \mu_6(e_1, e_4) = -e_6$ $\mu_6(e_2, e_3) = e_6$
\mathfrak{n}_7	$\mu_7^t(e_1, e_2) = -e_4, \mu_7^t(e_1, e_3) = \sqrt{t}e_5$ $\mu_7^t(e_2, e_4) = \sqrt{t}e_5, \mu_7^t(e_1, e_4) = -\frac{1}{\sqrt{t}}e_6$ $\mu_7^t(e_2, e_3) = \frac{1}{\sqrt{t}}e_6, t \in (0, 1]$
	$\tilde{\mu}_7^t(e_1, e_2) = -e_4, \tilde{\mu}_7^t(e_1, e_3) = \sqrt{-t}e_5$ $\tilde{\mu}_7^t(e_2, e_4) = \sqrt{-t}e_5, \tilde{\mu}_7^t(e_1, e_4) = \frac{1}{\sqrt{-t}}e_6$ $\tilde{\mu}_7^t(e_2, e_3) = -\frac{1}{\sqrt{-t}}e_6, t \in [-1, 0)$

Tab. 1. Abelian complex nilmanifolds of dimension 6.

Proof. We have

$$g = \begin{bmatrix} \frac{\sqrt{2}}{2}i & -\frac{\sqrt{2}}{2} & 0 \\ \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2}i & 0 \\ 0 & 0 & 1 \end{bmatrix} \in \mathrm{U}(2) \times \mathrm{U}(1).$$

Using the identification $a + bi \mapsto \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$, we thus get

$$(10) \quad g = \begin{bmatrix} 0 & -\frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 & 0 & 0 \\ \frac{\sqrt{2}}{2} & 0 & 0 & -\frac{\sqrt{2}}{2} & 0 & 0 \\ \frac{\sqrt{2}}{2} & 0 & 0 & \frac{\sqrt{2}}{2} & 0 & 0 \\ 0 & \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

By definition, it follows that

$$\begin{aligned}
& - \mu_4^{2,t}(e_1, e_2) = 0. \\
& g \cdot \mu_4^{1,t}(e_1, e_2) = g\mu_4^{1,t} \left(-\frac{\sqrt{2}}{2}e_2 - \frac{\sqrt{2}}{2}e_3, \frac{\sqrt{2}}{2}e_1 - \frac{\sqrt{2}}{2}e_4 \right) = g\left\{ \frac{1}{2} (\sqrt{t}e_5 - \sqrt{t}e_5) \right\} = 0. \\
& - \mu_4^{2,t}(e_1, e_3) = \sqrt{t}e_5. \\
& g \cdot \mu_4^{1,t}(e_1, e_3) = g\mu_4^{1,t} \left(-\frac{\sqrt{2}}{2}e_2 - \frac{\sqrt{2}}{2}e_3, \frac{\sqrt{2}}{2}e_1 + \frac{\sqrt{2}}{2}e_4 \right) = g\left\{ \frac{1}{2} (\sqrt{t}e_5 + \sqrt{t}e_5) \right\} \\
& = \sqrt{t}e_5. \\
& - \mu_4^{2,t}(e_1, e_4) = -\frac{1}{\sqrt{t}}e_6. \\
& g \cdot \mu_4^{1,t}(e_1, e_4) = g\mu_4^{1,t} \left(-\frac{\sqrt{2}}{2}e_2 - \frac{\sqrt{2}}{2}e_3, \frac{\sqrt{2}}{2}e_2 - \frac{\sqrt{2}}{2}e_3 \right) = g\left\{ \frac{1}{2} \left(-\frac{1}{\sqrt{t}}e_6 - \frac{1}{\sqrt{t}}e_6 \right) \right\} \\
& = -\frac{1}{\sqrt{t}}e_6. \\
& - \mu_4^{2,t}(e_2, e_3) = \frac{1}{\sqrt{t}}e_6. \\
& g \cdot \mu_4^{1,t}(e_2, e_3) = g\mu_4^{1,t} \left(\frac{\sqrt{2}}{2}e_1 - \frac{\sqrt{2}}{2}e_4, \frac{\sqrt{2}}{2}e_1 + \frac{\sqrt{2}}{2}e_4 \right) = g\left\{ \frac{1}{2} \left(\frac{1}{\sqrt{t}}e_6 + \frac{1}{\sqrt{t}}e_6 \right) \right\} \\
& = \frac{1}{\sqrt{t}}e_6. \\
& - \mu_4^{2,t}(e_2, e_4) = \sqrt{t}e_5. \\
& g \cdot \mu_4^{1,t}(e_2, e_4) = g\mu_4^{1,t} \left(\frac{\sqrt{2}}{2}e_1 - \frac{\sqrt{2}}{2}e_4, \frac{\sqrt{2}}{2}e_2 - \frac{\sqrt{2}}{2}e_3 \right) = g\left\{ \frac{1}{2} (\sqrt{t}e_5 + \sqrt{t}e_5) \right\} \\
& = \sqrt{t}e_5. \\
& - \mu_4^{2,t}(e_3, e_4) = 0. \\
& g \cdot \mu_4^{1,t}(e_3, e_4) = g\mu_4^{1,t} \left(\frac{\sqrt{2}}{2}e_1 + \frac{\sqrt{2}}{2}e_4, \frac{\sqrt{2}}{2}e_2 - \frac{\sqrt{2}}{2}e_3 \right) = g\left\{ \frac{1}{2} (\sqrt{t}e_5 - \sqrt{t}e_5) \right\} = 0.
\end{aligned}$$

Hence $g \cdot \mu_4^{1,t} = \mu_4^{2,t}$, which completes the proof. \square

Theorem 4.4. *Any 6-dimensional abelian complex nilmanifold admits a minimal metric, with the only exception of (N_5, J) .*

Proof. By applying Theorem 3.4 (as we described in Example 3.7 for \mathfrak{n}_4), it is easily seen that (N_1, J) admit a minimal metric of type $(3 < 5 < 6; 2, 2, 2)$; (N_2, J) , (N_3, J) and (N_4, J) one of type $(1 < 2; 4, 2)$; (N_6, J) and (N_7, J) one of type $(1 < 2 < 3; 2, 2, 2)$. Furthermore, we can see that each μ_i on \mathfrak{n}_i is minimal, if $i \neq 5$ (column 4, Table 2). Note that the Table 2 differs from the Table 1 in \mathfrak{n}_3 and \mathfrak{n}_7 , this is due to get μ_3 and μ_7 minimals was required to act with a matrix $g \in GL_3(\mathbb{C})$ in the brackets given in the Table 1. For example, for \mathfrak{n}_7 , take

$$g = \begin{bmatrix} \alpha & & \\ & \frac{1}{\alpha} & \\ & & 1 \end{bmatrix},$$

where $\alpha = (t + \frac{1}{t})^{-\frac{1}{6}}$ for μ_7^t , and $\alpha = (-t - \frac{1}{t})^{-\frac{1}{6}}$ for $\tilde{\mu}_7^t$.

It remains to prove that (N_5, J) does not admit minimal compatible metrics. To do this, we will use some properties of the $GL_n(\mathbb{R})$ -invariant stratification for the representation $\Lambda^2(\mathbb{R}^n)^* \otimes \mathbb{R}^n$ of $GL_n(\mathbb{R})$ (see [10], [9] for more details).

Let $\beta = \text{diag}(-1/2, -1/2, -1/2, -1/2, 1/2, 1/2)$. Hence

$$G_\beta := \{g \in GL(6) : g\beta g^{-1} = \beta, gJg^{-1} = J\} = GL_2(\mathbb{C}) \times GL_1(\mathbb{C})$$

n	Bracket	Type	Minimal
\mathfrak{n}_1	$\mu_1(e_1, e_2) = e_6$	$(3 < 5 < 6; 2, 2, 2)$	Yes
\mathfrak{n}_2	$\mu_2^\pm(e_1, e_2) = e_6, \mu_2^\pm(e_3, e_4) = \pm e_6$	$(1 < 2; 4, 2)$	Yes
\mathfrak{n}_3	$\mu_3^s(e_1, e_2) = e_5, \mu_3^s(e_3, e_4) = \frac{-s}{\sqrt{1+s^2}}e_5 + \frac{1}{\sqrt{1+s^2}}e_6$ $s \in \mathbb{R}$	$(1 < 2; 4, 2)$	Yes
\mathfrak{n}_4	$\mu_4^t(e_1, e_2) = \sqrt{t}e_5, \mu_4^t(e_1, e_4) = \frac{1}{\sqrt{t}}e_6$ $\mu_4^t(e_2, e_3) = -\frac{1}{\sqrt{t}}e_6, \mu_4^t(e_3, e_4) = -\sqrt{t}e_5$ $t \in (0, 1]$	$(1 < 2; 4, 2)$	Yes
\mathfrak{n}_5	$\mu_5(e_1, e_2) = e_5, \mu_5(e_1, e_4) = -e_6$ $\mu_5(e_2, e_3) = e_6$	—	No
\mathfrak{n}_6	$\mu_6(e_1, e_2) = -e_3, \mu_6(e_1, e_4) = -e_6$ $\mu_6(e_2, e_3) = e_6$	$(1 < 2 < 3; 2, 2, 2)$	Yes
\mathfrak{n}_7	$\mu_7^t(e_1, e_2) = -\sqrt{t+1}te_4, \mu_7^t(e_1, e_3) = \sqrt{t}e_5$ $\mu_7^t(e_2, e_4) = \sqrt{t}e_5, \mu_7^t(e_1, e_4) = -\frac{1}{\sqrt{t}}e_6$ $\mu_7^t(e_2, e_3) = \frac{1}{\sqrt{t}}e_6, t \in (0, 1]$	$(1 < 2 < 3; 2, 2, 2)$	Yes
	$\tilde{\mu}_7^t(e_1, e_2) = -\sqrt{-t-1}te_4, \tilde{\mu}_7^t(e_1, e_3) = \sqrt{-t}e_5$ $\tilde{\mu}_7^t(e_2, e_4) = \sqrt{-t}e_5, \tilde{\mu}_7^t(e_1, e_4) = \frac{1}{\sqrt{-t}}e_6$ $\tilde{\mu}_7^t(e_2, e_3) = -\frac{1}{\sqrt{-t}}e_6, t \in [-1, 0)$		

Tab. 2. Minimal metrics on 6-dimensional abelian complex nilmanifolds.

Since $\mathfrak{g}_\beta = \mathbb{R}\beta \oplus^\perp \mathfrak{h}_\beta$, it follows that \mathfrak{h}_β is Lie subalgebra. Let $H_\beta \subset G_\beta$ denote the Lie subgroup with Lie algebra \mathfrak{h}_β . We thus get

$$\mathfrak{h}_\beta = \left\{ \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} : \operatorname{tr} A = \operatorname{tr} B \right\}, \quad H_\beta = \left\{ \begin{bmatrix} g & 0 \\ 0 & h \end{bmatrix} : \det(g) = \det(h) \right\}.$$

But $\mathfrak{h}_\beta = (\mathbb{R} \begin{bmatrix} I & \\ & 2I \end{bmatrix}) \oplus \tilde{\mathfrak{h}}_\beta$ where

$$\tilde{\mathfrak{h}}_\beta = \left\{ \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} : \operatorname{tr} A = \operatorname{tr} B = 0 \right\}.$$

This clearly forces $\tilde{H}_\beta = \overline{SL_2(\mathbb{C})} \times \{I\}$. Therefore, it suffices to prove that $0 \notin \overline{SL_2(\mathbb{C})} \cdot \mu_5$ and $\mu_2 \in \overline{SL_2(\mathbb{C})} \cdot \mu_5$, with μ_2 and μ_5 the brackets of \mathfrak{n}_2 and \mathfrak{n}_5 respectively, which is due to the fact that $G \cdot \mu$ is minimal if and only if $H_\beta \cdot \mu$ is

closed (see for instance [9, Theorem 9.1]). Indeed, an easy computation shows that

$$\begin{bmatrix} a & & \\ & a & \\ & 1/a & \\ & & 1/a \end{bmatrix} \cdot \mu_5 \longrightarrow \mu_2 \quad \text{letting } a \rightarrow \infty.$$

From what has already been and the fact that $SL_2(\mathbb{C}) \cdot \mu_2$ is closed (\mathfrak{n}_2 is minimal), we conclude that $0 \notin \overline{SL_2(\mathbb{C}) \cdot \mu_5}$ by the uniqueness of closed orbits in the closure of an orbit (note that $\{0\}$ is a closed orbit). \square

We now will use the Pfaffian forms to give an alternative proof of the pairwise non-isomorphism of the family given in [1, Theorem 3.4] in the 2-step nilpotent case. Since $\dim \mathfrak{v}_1 = 4$ and $\dim \mathfrak{v}_2 = 2$, the Pfaffian forms of $\mathfrak{n}_1, \dots, \mathfrak{n}_5$ belong to the set $P_{2,2}(\mathbb{R})$; so we are left with the task of determining the quotient $P_{2,2}(\mathbb{R})/GL_1(\mathbb{C}) = P_{2,2}(\mathbb{R})/\mathbb{R}_{>0}U(1)$ (see Proposition 3.12).

Using the identification $P = ax^2 + bxy + cy^2 \leftrightarrow P_A := \langle A(x, y), (x, y) \rangle$, where $A = \begin{bmatrix} a & b/2 \\ b/2 & c \end{bmatrix}$, we have (see Remark 3.9)

$$P_{2,2}(\mathbb{R})/\pm GL_2(\mathbb{R}) = \begin{cases} x^2 + y^2, \\ x^2 - y^2, \\ x^2, \\ 0. \end{cases}$$

Proposition 3.12 now implies that

$$P_{2,2}(\mathbb{R})/\mathbb{R}_{>0}U(1) = \{ax^2 + by^2 : a \leq b, a^2 + b^2 = 1\} \cup \{0\}.$$

This allows us to classify the Pfaffian forms of $\mathfrak{n}_1, \dots, \mathfrak{n}_5$, which is summarized in Figure 1. The Lie algebra \mathfrak{n}_4^* is given by $\mu_t(e_1, e_3) = -tse_6$, $\mu_t(e_1, e_4) = \mu_t(e_2, e_3) = se_5$, $\mu_t(e_2, e_4) = s(2-t)e_6$, with $s = \sqrt{2+t^2 + (2-t)^2}$, $1 \leq t < 2$; it is minimal and (N_{μ_t}, J) is not abelian (see [7, Example 5.3]).

From Figure 1, it is clear that \mathfrak{n}_3 and \mathfrak{n}_4 have (minimal) Hermitian metric curves; $(\mathfrak{n}_2, \mu_2^+)$ and $(\mathfrak{n}_2, \mu_2^-)$ are distinguished; \mathfrak{n}_1 has an unique (minimal) Hermitian metric; and \mathfrak{n}_5 has an unique Hermitian metric.

We now consider the Lie algebras which are not 2-step nilpotent. The Lie algebra \mathfrak{n}_6 has an unique minimal metric up to isometry and scaling, by Theorem 3.4. For \mathfrak{n}_7 , an easy computation shows that for all $t \in [-1, 0)$, $s \in (0, 1]$

$$\text{Ric}_{\mu_7^t} \big|_{\mathfrak{z}} = \begin{bmatrix} -t & 0 \\ 0 & -1/t \end{bmatrix}, \quad \text{Ric}_{\mu_7^s} \big|_{\mathfrak{z}} = \begin{bmatrix} s & 0 \\ 0 & 1/s \end{bmatrix},$$

where $\mathfrak{z} := \langle e_5, e_6 \rangle_{\mathbb{R}}$. From this we deduce that the Hermitian nilmanifolds $\{(N_{\mu_7^t}, J, \langle \cdot, \cdot \rangle) : t \in (0, 1]\}$ are pairwise non-isometric (as we described in Example 3.7 for \mathfrak{n}_4). Likewise for $\{(N_{\mu_7^t}, J, \langle \cdot, \cdot \rangle) : t \in [-1, 0)\}$.

We will distinguish μ_7^t , $t \in (0, 1]$, of $\tilde{\mu}_7^t$, $t \in [-1, 0)$. To do this we need the following (see (2))

$$\begin{aligned} \|\mu_7^t\|^2 &= 2(\|\mu_7^t(e_1, e_2)\|^2 + \|\mu_7^t(e_1, e_3)\|^2 + \|\mu_7^t(e_1, e_4)\|^2 + \|\mu_7^t(e_2, e_3)\|^2 \\ &\quad + \|\mu_7^t(e_2, e_4)\|^2) = 6\left(t + \frac{1}{t}\right), \quad t \in (0, 1]. \end{aligned}$$

$$\begin{aligned} \|\tilde{\mu}_7^t\|^2 &= 2(\|\tilde{\mu}_7^t(e_1, e_2)\|^2 + \|\tilde{\mu}_7^t(e_1, e_3)\|^2 + \|\tilde{\mu}_7^t(e_1, e_4)\|^2 + \|\tilde{\mu}_7^t(e_2, e_3)\|^2 \\ &\quad + \|\tilde{\mu}_7^t(e_2, e_4)\|^2) = -6\left(t + \frac{1}{t}\right), \quad t \in [-1, 0). \end{aligned}$$




Fig. 1. Pfaffian forms of $\mathbf{n}_1, \dots, \mathbf{n}_5$.

Proposition 4.5. $\tilde{\mu}_7^t \notin \mathbb{R}^* \text{U}(1) \times \text{U}(1) \times \text{U}(1) \cdot \mu_7^s$ for all $t \in [-1, 0)$, $s \in (0, 1]$.

Proof. If we suppose that there exists $c \in \mathbb{R}^*$ and $\varphi \in \text{U}(1) \times \text{U}(1) \times \text{U}(1)$ such that $c\tilde{\mu}_7^t = \varphi \cdot \mu_7^s$, then $\varphi = \begin{bmatrix} \varphi_1 & & \\ & \varphi_2 & \\ & & \varphi_3 \end{bmatrix}$ and $c^2 \text{Ric}_{\tilde{\mu}_7^t}|_3 = \varphi_3 \text{Ric}_{\mu_7^s}|_3 \varphi_3^{-1}$. Hence $c^2 \begin{bmatrix} -t & \\ & -1/t \end{bmatrix} = \begin{bmatrix} s & \\ & 1/s \end{bmatrix}$; taking quotients of their eigenvalues we deduce that $s^2 = t^2$ or $s^2 = 1/t^2$, which gives $t = -s$ if $t \in [-1, 0)$, $s \in (0, 1]$. From this it is enough to prove that for all $t \in (0, 1]$, $c \in \mathbb{R}^*$,

$$(11) \quad \tilde{\mu}_7^{-t} \notin c \text{U}(1) \times \text{U}(1) \times \text{U}(1) \cdot \mu_7^t.$$

Moreover, if $\tilde{\mu}_7^{-t} \in c \text{U}(1) \times \text{U}(1) \times \text{U}(1) \cdot \mu_7^t$, then $\|\tilde{\mu}_7^{-t}\|^2 = c^2 \|\mu_7^t\|^2$, which yields $c^2 = 1$, and hence $c = \pm 1$. Thus it is sufficient to take $c = 1$ (if $c = -1$ the equations does not change).

Suppose, contrary to our claim, that $\tilde{\mu}_7^{-t} = G \cdot \mu_7^t$ where

$$G = \begin{bmatrix} a & -b & & & & \\ b & a & & & & \\ & & c & -d & & \\ & & d & c & & \\ & & & & k & -h \\ & & & & h & k \end{bmatrix}, \quad G^{-1} = \begin{bmatrix} a & b & & & & \\ -b & a & & & & \\ & & c & d & & \\ & & -d & c & & \\ & & & & k & h \\ & & & & -h & k \end{bmatrix},$$

with $a^2 + b^2 = c^2 + d^2 = k^2 + h^2 = 1$. We thus get

- $\tilde{\mu}_7^{-t}(e_1, e_2) = -\sqrt{t+1}/te_4 = G \cdot \mu_7^t(e_1, e_2) = d\sqrt{t+1}/te_3 - c\sqrt{t+1}/te_4.$
- $\tilde{\mu}_7^{-t}(e_1, e_3) = \sqrt{t}e_5 = G \cdot \mu_7^t(e_1, e_3) = \{(ac + bd)k\sqrt{t} + (bc - ad)\frac{h}{\sqrt{t}}\}e_5$
 $+ \{(ac + bd)h\sqrt{t} + (ad - bc)\frac{k}{\sqrt{t}}\}e_6.$
- $\tilde{\mu}_7^{-t}(e_1, e_4) = \frac{1}{\sqrt{t}}e_6 = G \cdot \mu_7^t(e_1, e_4) = \{(ad - bc)k\sqrt{t} + (ac + bd)\frac{h}{\sqrt{t}}\}e_5$
 $+ \{(ad - bc)h\sqrt{t} - (ac + bd)\frac{k}{\sqrt{t}}\}e_6.$

This is equivalent at next system (the other tree brackets produce the same equations):

$$\begin{cases} c = 1, & d = 0, \\ a = k, \\ b - ht = 0, \\ a = -k, \\ h + bt = 0, \end{cases}$$

It follows easily that $a = b = 0$, contrary to $a^2 + b^2 = 1$. Since G was arbitrary, (11) is proved. \square

5. RESULTS OBTAINED IN DIMENSION EIGHT

In this section, our aim is to exhibit many families depending on one (see Example 5.4 and Example 5.8), two (see Example 5.3 and Example 5.5) and three (see Example 5.2) parameters of abelian complex structures on 8-dimensional 2-step nilpotent Lie algebras, by using that they all admit minimal metrics for the types $(1 < 2; 4, 4)$ and $(1 < 2; 6, 2)$.

Following the idea developed in dimension six, we will determine the quotients $P_{4,2}(\mathbb{R})/\mathbb{R}_{>0}\mathrm{U}(2)$ and $P_{2,3}(\mathbb{R})/\mathbb{R}_{>0}\mathrm{U}(1)$ in the cases $(4, 4)$ and $(6, 2)$ respectively. This may be viewed as a first step towards the classification of abelian complex structures on 8-dimensional nilmanifolds. From now on, we keep the notation used in [6].

5.1. Type $(4,4)$.

In this case $\mathfrak{v}_1 = \mathbb{R}^4$ and $\mathfrak{v}_2 = \mathbb{R}^4$, and we consider the vector space $W := \Lambda^2 \mathfrak{v}_1^* \otimes \mathfrak{v}_2$. If $\{X_1, \dots, X_4, Z_1, \dots, Z_4\}$ is a basis of \mathfrak{n} such that $\mathfrak{v}_1 = \langle X_1, \dots, X_4 \rangle_{\mathbb{R}}$ and $\mathfrak{v}_2 = \langle Z_1, \dots, Z_4 \rangle_{\mathbb{R}}$, then each element in W will be described as

$$\begin{aligned} \mu(X_1, X_2) &= a_1 Z_1 + a_2 Z_2 + a_3 Z_3 + a_4 Z_4, & \mu(X_1, X_3) &= b_1 Z_1 + b_2 Z_2 + b_3 Z_3 + b_4 Z_4, \\ \mu(X_1, X_4) &= c_1 Z_1 + c_2 Z_2 + c_3 Z_3 + c_4 Z_4, & \mu(X_2, X_3) &= d_1 Z_1 + d_2 Z_2 + d_3 Z_3 + d_4 Z_4, \\ \mu(X_2, X_4) &= e_1 Z_1 + e_2 Z_2 + e_3 Z_3 + e_4 Z_4, & \mu(X_3, X_4) &= f_1 Z_1 + f_2 Z_2 + f_3 Z_3 + f_4 Z_4. \end{aligned}$$

The complex structure and the compatible metric will be always defined by

$$J = \begin{bmatrix} 0 & -1 & & & & \\ 1 & 0 & & & & \\ & & 0 & -1 & & \\ & & 1 & 0 & & \\ & & & & 0 & -1 \\ & & & & 1 & 0 \\ & & & & & & 0 & -1 \\ & & & & & & 1 & 0 \end{bmatrix}, \quad \langle X_i, X_j \rangle = \langle Z_i, Z_j \rangle = \delta_{ij}.$$

If $A = (a_1, \dots, a_4), \dots, F = (f_1, \dots, f_4)$ and $JA = (-a_2, a_1, -a_4, a_3), \dots, JF = (-f_2, f_1, -f_4, f_3)$, then J is integrable on N_μ (i.e. J satisfies (1)), $\mu \in W$, if and only if

$$(12) \quad E = B + JC + JD,$$

and J is abelian if and only if

$$(13) \quad E = B, \quad D = -C.$$

Define $v_i = (a_i, b_i, c_i, d_i, e_i, f_i)$, $i = 1, 2, 3, 4$. It is easy to check that for any $\mu \in W$, $\text{Ric}_\mu|_{v_2} = \frac{1}{2}[\langle v_i, v_j \rangle]$, $1 \leq i, j \leq 4$, and

$$\text{Ric}_\mu|_{v_1} = -\frac{1}{2} \begin{bmatrix} \|A\|^2 + \|B\|^2 + \|C\|^2 & \langle B, D \rangle + \langle C, E \rangle & -\langle A, D \rangle + \langle C, F \rangle & -\langle A, E \rangle - \langle B, F \rangle \\ \langle B, D \rangle + \langle C, E \rangle & \|A\|^2 + \|D\|^2 + \|E\|^2 & \langle A, B \rangle + \langle E, F \rangle & \langle A, C \rangle - \langle D, F \rangle \\ -\langle A, D \rangle + \langle C, F \rangle & \langle A, B \rangle + \langle E, F \rangle & \|B\|^2 + \|D\|^2 + \|F\|^2 & \langle B, C \rangle + \langle D, E \rangle \\ -\langle A, E \rangle - \langle B, F \rangle & \langle A, C \rangle - \langle D, F \rangle & \langle B, C \rangle + \langle D, E \rangle & \|C\|^2 + \|E\|^2 + \|F\|^2 \end{bmatrix}.$$

Therefore

$$\begin{aligned} & \text{Ric}_\mu^c|_{v_1} \\ &= \frac{1}{4} \begin{bmatrix} -\alpha & 0 & \langle A + F, D - C \rangle & \langle A + F, B + E \rangle \\ 0 & -\alpha & -\langle A + F, B + E \rangle & \langle A + F, D - C \rangle \\ \langle A + F, D - C \rangle & -\langle A + F, B + E \rangle & -\beta & 0 \\ -\langle A + F, B + E \rangle & \langle A + F, D - C \rangle & 0 & -\beta \end{bmatrix}, \end{aligned}$$

$$\begin{aligned} & \text{Ric}_\mu^c|_{v_2} \\ &= \frac{1}{4} \begin{bmatrix} \|v_1\|^2 + \|v_2\|^2 & 0 & \langle v_1, v_3 \rangle + \langle v_2, v_4 \rangle & \langle v_1, v_4 \rangle - \langle v_2, v_3 \rangle \\ 0 & \|v_1\|^2 + \|v_2\|^2 & \langle v_2, v_3 \rangle - \langle v_2, v_4 \rangle & \langle v_2, v_4 \rangle + \langle v_1, v_3 \rangle \\ \langle v_1, v_3 \rangle + \langle v_2, v_4 \rangle & \langle v_2, v_3 \rangle - \langle v_2, v_4 \rangle & \|v_3\|^2 + \|v_4\|^2 & 0 \\ \langle v_1, v_4 \rangle - \langle v_2, v_3 \rangle & \langle v_2, v_4 \rangle + \langle v_1, v_3 \rangle & 0 & \|v_3\|^2 + \|v_4\|^2 \end{bmatrix}, \end{aligned}$$

where $\alpha := 2\|A\|^2 + \|B\|^2 + \|C\|^2 + \|D\|^2 + \|E\|^2$ and $\beta := \|B\|^2 + \|C\|^2 + \|D\|^2 + \|E\|^2 + 2\|F\|^2$.

One type of minimality which is easy to characterize is $(1 < 2; 4, 4)$. Indeed, if for any $\mu \in W$ we have that $\text{Ric}_\mu^c|_{v_1} = pI_4$ and $\text{Ric}_\mu^c|_{v_2} = qI_4$, then

$$\text{Ric}_\mu^c = \begin{bmatrix} pI_4 & \\ & qI_4 \end{bmatrix} = (2p - q)I_8 + (q - p) \begin{bmatrix} I_4 & \\ & 2I_4 \end{bmatrix} \in \mathbb{R}I + \text{Der}(\mu).$$

The following are sufficient conditions for any $\mu \in W$ is minimal of type $(1 < 2; 4, 4)$.

(i) Conditions for $\text{Ric}_\mu^c|_{n_1} \in \mathbb{R}I$:

- $\langle A + F, D - C \rangle = 0$.
- $\langle A + F, B + E \rangle = 0$.

- $\|A\|^2 = \|F\|^2$.

(ii) Conditions for $\text{Ric}^c_\mu|_{\mathfrak{n}_2} \in \mathbb{R}I$:

- $\|v_1\|^2 + \|v_2\|^2 = \|v_3\|^2 + \|v_4\|^2$.
- $\langle v_1, v_3 \rangle = -\langle v_2, v_4 \rangle$.
- $\langle v_1, v_4 \rangle = \langle v_2, v_3 \rangle$.

Moreover, if μ satisfies the conditions given in (i) and (ii), we obtain $p = -\frac{1}{4}\alpha$ and $q = \frac{1}{4}(\|v_1\|^2 + \|v_2\|^2)$.

In the rest of this section we will study the Pfaffian forms of $\mu \in W$. Since $\dim \mathfrak{v}_1 = \dim \mathfrak{v}_2 = 4$, it follows that the Pfaffian form of any $\mu \in W$ belongs to the set $P_{4,2}(\mathbb{R})$; so the goal is to determine the quotient $P_{4,2}(\mathbb{R})/\mathbb{R}_{>0}U(2)$. As in the case (4, 2) there is the identification $f(\mu) \in P_{4,2}(\mathbb{R}) \leftrightarrow A_f$, where A_f is a symmetric matrix, and, in consequence,

$$(14) \quad P_{4,2}(\mathbb{R})/\pm GL_4(\mathbb{R}) = \left\{ \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & 0 \end{bmatrix}, \dots \right\}.$$

Based on the classification of complex metabelian (two-step nilpotent) Lie algebras in dimension up to 9 given by L. Yu. Galitski and D. A. Timashev in [5], and by using the identifications of the real forms of Lie algebra on \mathbb{C} , we have

$$(15) \quad P_{4,2}(\mathbb{C})/GL_2(\mathbb{C}) = \begin{cases} x^2 - y^2 - z^2 + w^2 \\ x^2 - y^2 - z^2 \\ x^2 - y^2 \\ x^2 \\ 0 \end{cases} \simeq \begin{cases} + & + & + & + \\ + & + & + & 0 \\ + & + & 0 & 0 \\ + & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{cases}$$

Remark 5.1. The polynomial $f = x^2 + y^2 + z^2 + w^2$ is not the Pfaffian form of any $\mu \in W$. In general, $f > 0$ ($\Leftrightarrow J_Z$ are invertible $\forall Z$) is not the Pfaffian form of any $\mu \in W$. The dimensions allowed for this are: $(2k, 1)$, $(4k, 2)$, $(4k, 3)$, $(8k, 4)$, \dots , $(8k, 7)$, $(16k, 8)$, $(32k, 9)$.

The following expression was obtained by direct calculation rather than the equations (14) and (15).

$$\begin{aligned}
P_{4,2}(\mathbb{R})/U(2) &\simeq \text{sym}(4)/U(2) \\
&= \begin{cases} \left(aI, \begin{bmatrix} b & & & \\ & -b & & \\ & & c & \\ & & & -c \end{bmatrix} \right); & a, b, c \in \mathbb{R}. \\ \\ \left(\begin{bmatrix} a & & & \\ & a & & \\ & & b & \\ & & & b \end{bmatrix}, \begin{bmatrix} c & h & & \\ h & -c & & \\ & & d & l \\ & & & -d \end{bmatrix} \right); & a, b, c, d, h, l \in \mathbb{R} \ (a < b). \end{cases} \\
&= \begin{cases} \begin{cases} ax^2 + by^2 + cz^2 + dw^2; & a + b = c + d, \\ a, b, c, d \in \mathbb{R}. \end{cases} \\ \\ \begin{cases} ax^2 + by^2 + cz^2 + dw^2 + hxy + lzw; & a + b < c + d, \\ a, b, c, d, h, l \in \mathbb{R}. \end{cases} \end{cases}
\end{aligned}$$

In what follows, we give some curves and families of minimal metrics of type $(1 < 2; 4, 4)$, which Pfaffian forms appear in the above quotient.

Example 5.2. Let $\mu_{krst} \in W$ be given by

$$\begin{aligned}
A &= (s, t, 0, 0), & B &= (0, 0, r, 0), \\
C &= (0, 0, 0, k), & D &= (0, 0, 0, -k), \\
E &= (0, 0, r, 0), & F &= (s, -t, 0, 0),
\end{aligned}$$

with $k, r, s, t \in \mathbb{R}$. It is clear that μ_{krst} satisfies (12) and (13), and hence $(N_{\mu_{krst}}, J)$ is an abelian complex nilmanifold for all $k, r, s, t \in \mathbb{R}$. Furthermore, if $k^2 + r^2 = s^2 + t^2$ then the family $\{(N_{\mu_{krst}}, J, \langle \cdot, \cdot \rangle) : k^2 + r^2 = s^2 + t^2\}$ of minimal (conditions (i) and (ii)) metrics is pairwise non-isometric, up to scaling. This gives rise then a 3-parameter family of pairwise non-isomorphic abelian complex nilpotent Lie groups (see Theorem 3.4). On the other hand, the Pfaffian form of μ_{krst} is

$$f(\mu_{krst}) = s^2x^2 - t^2y^2 - r^2z^2 - k^2w^2.$$

Example 5.3. Let λ_{rst} be defined by:

$$\begin{aligned}
A &= (0, r, 0, 0), & B &= (0, 0, s, 0), \\
C &= (0, 0, 0, t), & D &= (0, 0, 0, -t), \\
E &= (0, 0, s, 0), & F &= (0, -r, 0, 0),
\end{aligned}$$

where $r, s, t \in \mathbb{R}$. We have $(N_{\lambda_{rst}}, J)$ is an abelian complex nilmanifold for all r, s, t in \mathbb{R} . If $r^2 = s^2 + t^2$ then the family $\{(N_{\lambda_{rst}}, J, \langle \cdot, \cdot \rangle) : r^2 = s^2 + t^2\}$ of minimal compatible metrics is pairwise non-isometric, unless scalar multiples. This gives rise then a 2-parameter family of pairwise non-isomorphic abelian complex nilpotent Lie groups. Note that the Pfaffian form of λ_{rst} is

$$f(\lambda_{rst}) = -r^2y^2 - s^2z^2 - t^2w^2.$$

Example 5.4. Let ν_{st} be given by $A = (s, 0, 0, 0) = -F$, $B = E = 0$, $C = (0, 0, t, 0) = -D$, with $s, t \in \mathbb{R}$. Therefore, $(N_{\nu_{st}}, J)$ is an abelian complex nilmanifold for all $s, t \in \mathbb{R}$. Furthermore, if $s^2 = t^2$ then the curve $\{\nu_{st} : s^2 = t^2\}$ of minimal compatible metrics is pairwise non-isometric, unless scalar multiples. This gives a curve of pairwise non-isomorphic abelian complex nilpotent Lie groups. Finally, the Pfaffian form of ν_{st} is

$$f(\nu_{st}) = -s^2x^2 - t^2z^2.$$

Example 5.5. Let $\mu_{rst} \in W$ be defined by:

$$\begin{aligned} A &= (r, 0, 0, 0), & B &= (0, 0, s, 0), \\ C &= (0, 0, 0, t), & D &= (0, 0, 0, -t), \\ E &= (0, 0, s, 0), & F &= (0, r, 0, 0), \end{aligned}$$

where $r, s, t \in \mathbb{R}$. Hence $(N_{\mu_{rst}}, J)$ is an abelian complex nilmanifold for all $r, s, t \in \mathbb{R}$. If $r^2 = s^2 + t^2$ then $\{\mu_{rst} : r^2 = s^2 + t^2\}$ of minimal compatible metrics is pairwise non-isometric, up to scaling. This gives rise then a 2-parameter family of pairwise non-isomorphic abelian complex nilpotent Lie groups. Note that the Pfaffian form of μ_{rst} is given by

$$f(\mu_{rst}) = r^2xy - s^2z^2 - t^2w^2.$$

5.2. Type (6,2).

For $\mathfrak{v}_1 = \mathbb{R}^6$ and $\mathfrak{v}_2 = \mathbb{R}^2$, consider $\widetilde{W} := \Lambda^2 \mathfrak{v}_1^* \otimes \mathfrak{v}_2$. Fix basis $\{X_1, \dots, X_6\}$ and $\{Z_1, Z_2\}$ of \mathfrak{v}_1 and \mathfrak{v}_2 , respectively. Each element $\mu \in \widetilde{W}$ will be described as

$$\begin{aligned} \mu(X_1, X_2) &= a_1Z_1 + a_2Z_2, & \mu(X_1, X_3) &= b_1Z_1 + b_2Z_2, & \mu(X_1, X_4) &= c_1Z_1 + c_2Z_2, \\ \mu(X_1, X_5) &= d_1Z_1 + d_2Z_2, & \mu(X_1, X_6) &= e_1Z_1 + e_2Z_2, & \mu(X_2, X_3) &= f_1Z_1 + f_2Z_2, \\ \mu(X_2, X_4) &= g_1Z_1 + g_2Z_2, & \mu(X_2, X_5) &= h_1Z_1 + h_2Z_2, & \mu(X_2, X_6) &= i_1Z_1 + i_2Z_2, \\ \mu(X_3, X_4) &= k_1Z_1 + k_2Z_2, & \mu(X_3, X_5) &= l_1Z_1 + l_2Z_2, & \mu(X_3, X_6) &= m_1Z_1 + m_2Z_2, \\ \mu(X_4, X_5) &= n_1Z_1 + n_2Z_2, & \mu(X_4, X_6) &= p_1Z_1 + p_2Z_2, & \mu(X_5, X_6) &= q_1Z_1 + q_2Z_2. \end{aligned}$$

The complex structure and the compatible metric will be always defined by

$$\begin{aligned} JX_1 &= X_2, & JX_3 &= X_4, & \langle X_i, X_j \rangle &= \delta_{ij}, & \langle Z_k, Z_l \rangle &= \delta_{kl}. \\ JX_5 &= X_6, & JZ_1 &= Z_2. \end{aligned}$$

If $A = (a_1, a_2), \dots, Q = (q_1, q_2)$ and $JA = (-a_2, a_1), \dots, JQ = (-q_2, q_1)$, then J satisfies (1) if and only if

$$(16) \quad G = B + JC + JF, \quad I = D + JE + JH, \quad P = L + JM + JN,$$

and J is abelian if and only if

$$(17) \quad B = G, \quad C = -F, \quad D = I, \quad E = -H, \quad L = P, \quad M = N.$$

Let $v_i = (a_i, b_i, c_i, d_i, e_i, f_i)$, $i = 1, 2$. It follows that $\text{Ric}_\mu|_{\mathfrak{v}_2} = \frac{1}{2}[\langle v_i, v_j \rangle]$, $1 \leq i, j \leq 2$, and

$$\text{Ric}^c_\mu|_{\mathfrak{v}_2} = \frac{1}{4} \begin{bmatrix} \|v_1\|^2 + \|v_2\|^2 & 0 \\ 0 & \|v_1\|^2 + \|v_2\|^2 \end{bmatrix} \in \mathbb{R}I.$$

For the complicated expressions, we only give sufficient conditions for any $\mu \in \widetilde{W}$ is minimal of type $(1 < 2; 6, 2)$ when J is abelian.

(†) Conditions for $\text{Ric}^c_\mu|_{\mathfrak{n}_1} \in \mathbb{R}I$:

- (a) $\|A\|^2 = \|K\|^2 = \|Q\|^2$, $\|B\|^2 + \|C\|^2 = \|D\|^2 + \|E\|^2 = \|L\|^2 + \|M\|^2$.
- (b) $\langle A + K, B \rangle = \langle A + K, C \rangle = \langle A + Q, D \rangle = \langle A + Q, E \rangle = \langle K + Q, L \rangle = \langle K + Q, M \rangle = 0$.
- (c) $\langle B, L \rangle = -\langle C, N \rangle$, $\langle B, M \rangle = -\langle C, P \rangle$, $\langle B, D \rangle = -\langle C, E \rangle$.
- (d) $\langle C, D \rangle = -\langle G, H \rangle$, $\langle D, L \rangle = -\langle E, M \rangle$, $\langle H, L \rangle = -\langle I, M \rangle$.

If μ satisfies the conditions given in (†) we thus get $q := \frac{1}{4} (\|v_1\|^2 + \|v_2\|^2)$ and $p := -\frac{1}{2} (\|A\|^2 + \|B\|^2 + \|C\|^2 + \|D\|^2 + \|E\|^2)$.

Since $\dim \mathfrak{v}_1 = 6$ and $\dim \mathfrak{v}_2 = 2$, the Pfaffian form of any $\mu \in \widetilde{W}$ belongs to the set $P_{2,3}(\mathbb{R})$. Unlike the previous two cases, there is no identification of $f(\mu) \in P_{2,3}(\mathbb{R})$ with a matrix, but it is known that every polynomial in $P_{2,3}(\mathbb{R})$ is the Pfaffian form of some $\mu \in \widetilde{W}$ (see [8]). Again, of [5], we obtain

$$(18) \quad P_{2,3}(\mathbb{C})/GL_2(\mathbb{C}) = \begin{cases} x^3 \\ x^2y + xy^2 = xy(x+y) \\ x^3 + x^2y = x^2(x+y) \simeq x^2y \end{cases}$$

But it is easy to see that

$$\begin{aligned} P_{2,3}(\mathbb{R}) \cap GL_2(\mathbb{C}) \cdot x^3 &= GL_2(\mathbb{R}) \cdot x^3, \\ P_{2,3}(\mathbb{R}) \cap GL_2(\mathbb{C}) \cdot (x^2y + xy^2) &= GL_2(\mathbb{R}) \cdot (x^2y + xy^2), \\ P_{2,3}(\mathbb{R}) \cap GL_2(\mathbb{C}) \cdot x^2y &= GL_2(\mathbb{R}) \cdot x^2y, \end{aligned}$$

and therefore

$$P_{2,3}(\mathbb{R})/GL_2(\mathbb{R}) = \begin{cases} x^3 \\ x^2y + xy^2 = xy(x+y) \\ x^3 + x^2y = x^2(x+y) \simeq x^2y \end{cases}$$

Example 5.6. Let $\mu_{st}^1, \mu_{st}^2, \mu_{st}^3 \in \widetilde{W}$ be defined by: for all $s, t \in \mathbb{R}$,

$$\begin{aligned} A^1 &= (0, s), & A^2 &= (t, 0), & A^3 &= (0, s), \\ E^1 &= (t, 0), & K^2 &= (0, s), & E^3 &= (t, 0), \\ H^1 &= (-t, 0), & Q^2 &= (s, t), & H^3 &= (-t, 0), \\ K^1 &= (s, 0), & & & K^3 &= (s, 0), \\ & & & & Q^3 &= (-s, 0). \end{aligned}$$

It follows immediately that $(N_{\mu_{st}^1}, J)$, $(N_{\mu_{st}^2}, J)$ and $(N_{\mu_{st}^3}, J)$ are abelian complex nilmanifolds for all $s, t \in \mathbb{R}$, as they satisfy (16) and (17). Furthermore, they are

not minimal of type $(1 < 2; 6, 2)$ and its Pfaffian forms are given by

$$f(\mu_{st}^1) = st^2x^3, \quad f(\mu_{st}^2) = s^2tx^2y + st^2xy^2, \quad f(\mu_{st}^3) = st^2x^3 + s^3x^2y.$$

Hence $\{(N_{\mu_{st}^2}, J) : s, t \in \mathbb{R} \setminus \{0, \pm 1\}\}$ and $\{(N_{\mu_{st}^3}, J) : s, t \in \mathbb{R} \setminus \{0\}\}$ are curves of abelian complex nilmanifolds, which is due to the fact that

$$\begin{aligned} \forall a, b \in \mathbb{R} \setminus \{0, \pm 1\}, a \neq b : x^2y + axy^2 &\notin \text{U}(1) \cdot (x^2y + bxy^2). \\ \forall a, b \in \mathbb{R} \setminus \{0\}, a \neq b : x^3 + ax^2y &\notin \text{U}(1) \cdot (x^3 + bx^2y). \end{aligned}$$

Remark 5.7. Let $p(x, y) = \sum_{i=0}^3 a_i x^{3-i} y^i \in P_{2,3}(\mathbb{R})$. Define

$$\begin{aligned} \Delta(p) &:= (3a_0 + a_2)^2 + (a_1 + 3a_3)^2. \\ \|p\|^2 &:= 6a_0^2 + 2a_1^2 + 2a_2^2 + 6a_3^2. \\ D(p) &:= 18a_0a_1a_2a_3 + a_1^2a_2^2 - 4a_0a_2^3 - 4a_1^3a_3 - 27a_0^2a_3^2. \end{aligned}$$

We have Δ is $SO(2)$ -invariant, $\|\cdot\|^2$ is $O(2)$ -invariant and D is $SL_2(\mathbb{R})$ -invariant.

Note that using quotients of the above invariants we can also obtain that $\{(N_{\mu_{st}^2}, J) : s, t \in \mathbb{R} \setminus \{0, \pm 1\}\}$ and $\{(N_{\mu_{st}^3}, J) : s, t \in \mathbb{R} \setminus \{0\}\}$ are curves of abelian complex nilmanifolds.

Example 5.8. Let $\lambda_{st} \in \widetilde{W}$ be given by $A = (t, s)$, $K = (-s, t)$ and $Q = (s, t)$, with $s, t \in \mathbb{R}$. We obtain $(N_{\lambda_{st}}, J)$ is an abelian complex nilmanifold for all $s, t \in \mathbb{R}$. Furthermore, λ_{st} is minimal of type $(1 < 2; 6, 2)$. On the other hand, the Pfaffian form of λ_{st} is

$$f(\lambda_{st}) = s^2tx^3 + s^3x^2y - t^3xy^2 - st^2y^3.$$

Define $a := s^2$, $b := t^2$, and consider

$$h(a, b) := \frac{D(f(\lambda_{st}))}{(\Delta(f(\lambda_{st})))^2} = \frac{4ab(a^2 - b^2)^2}{(a + b)^6}.$$

If $a + b = 1$ then $h(a) = 4a(1 - a)(2a - 1)^2$ is an injective function for all $a \geq 1$. Hence $\{\lambda_{st} : s^2 + t^2 = 1, s \geq 1\}$ is a curve of pairwise non-isometric metrics.

REFERENCES

- [1] Andrada, A., Barberis, M.L., Dotti, I.G., *Classification of abelian complex structures on 6-dimensional Lie algebras*, J. London Math. Soc. **83** (1) (2011), 232–255.
- [2] Ceballos, M., Otal, A., Ugarte, L., Villacampa, R., *Classification of complex structures on 6-dimensional nilpotent Lie algebras*, arXiv:math.DG/1111.5873.
- [3] Cordero, L.A., Fernández, M., Ugarte, L., *Abelian complex structures on 6-dimensional compact nilmanifolds*, Comment. Math. Univ. Carolin. **43** (2) (2002), 215–229.
- [4] Dolgachev, I., *Lectures on invariant theory*, London Math. Soc. Lecture Note Ser. **296** (2003), 1–220.
- [5] Galitski, L.Yu., Timashev, D.A., *On classification of metabelian Lie algebras*, J. Lie Theory **9** (1999), 125–156.
- [6] Lauret, J., *Minimal metrics on nilmanifolds*, Differential geometry and its applications, Matfyzpress, Prague, 2005, pp. 79–97.

- [7] Lauret, J., *A canonical compatible metric for geometric structures on nilmanifolds*, Ann. Global Anal. Geom. **30** (2006), 107–138.
- [8] Lauret, J., *Rational forms of nilpotent Lie algebras and Anosov diffeomorphisms*, Monatsh. Math. **155** (2008), 15–30.
- [9] Lauret, J., *Einstein solvmanifolds and nilsolitons*, Contemp. Math. **491** (2009), 1–35.
- [10] Lauret, J., *Einstein solvmanifolds are standard*, Ann. of Math. (2) **172** (2010), 1859–1877.
- [11] Salamon, S.M., *Complex structures on nilpotent Lie algebras*, J. Pure Appl. Algebra **157** (2001), 311–333.

FAMAF AND CIEM, UNIVERSIDAD NACIONAL DE CÓRDOBA,
MEDINA ALLENDE S/N,
5000 CÓRDOBA, ARGENTINA
E-mail: earodriguez@famaf.unc.edu.ar